US Family Health Plan Prior Authorization Request Form for

Androgel, testosterone 1% and 1.62% gel MDP and gel packets, 2% testosterone solution MDP, 1% testosterone gel in unit-dose tubes (Testim, Vogelxo, generics)

To be completed and signed by the prescriber. To be used only for prescriptions which are to be filled through the Department of Defense (DoD) US Family Health Plan Pharmacy Program. US Family Health Plan is a TRICARE contractor for DoD.

The completed form may be faxed to 855-273-5735

OR

The patient may attach the completed form to the prescription and **mail** it to:

Attn: Pharmacy, 77 Warren St, Brighton, MA 02135

QUESTIONS? Call 1-877-880-7007

https://www.usfamilyhealth.org/for-providers/pharmacy-information/

Initial therapy approves for 1 year, renewal approves indefinitely. For renewal of therapy, an initial Tricare prior authorization approval is required. Clinical documentation including lab work may be required.

| Step | Please complete patient and physician information (please print): | | | | | |
|------|--|---|-----------------------------------|-----------------------|--|--|
| 1 | Patient Name: Address: Sponsor ID # Date of Birth: | | Physician Name: | | | |
| | | | Address: | | | |
| | | | Dhana th | | | |
| | | | Phone #: Secure Fax #: | | | |
| Step | Please complete the clinical assessment: | | | | | |
| 2 | A MEN di a conservati di a con | | | | | |
| | Will the requested medication be used to enhance athletic performance? | | ☐ Yes | □ No | | |
| | | | STOP | Proceed to question 2 | | |
| | | | Coverage not approved | | | |
| | 2. | Will the requested medication be used concomitantly with other testosterone products? | Yes | No | | |
| | | | STOP | Proceed to question 3 | | |
| | | | Coverage not approved | | | |
| | 3. | | ☐ Yes | □ No | | |
| | | under the TRICARE benefit in the last 6 months? Please choose "No" if the patient did not previously have a TRICARE approved PA for the requested medication. | Proceed to question 4 | Proceed to question 8 | | |
| | 4. | What is the indication or diagnosis? | ☐ Hypogonadism - Proceed to quest | ion 6 | | |
| | | ☐ Female-to-male gender dysphoria hormone therapy in a natal female patient (assigned female at birth) - Proceed to question 5 | | | | |
| | | | ☐ Other - Proceed to question 6 | | | |
| | 5. | Is the patient 19 years of age or older? | □ Yes | □ No | | |
| | | | Proceed to question 6 | STOP | | |
| | | | | Coverage not approved | | |
| | 6. | Has the patient had a positive response to | □ Yes | □ No | | |
| | | therapy? | Proceed to question 7 | STOP | | |
| | | | | Coverage not approved | | |

Tricare Prior Authorization Request Form for

Androgel, testosterone 1% and 1.62% gel MDP and gel packets, 2% testosterone solution MDP, 1% testosterone gel in unit-dose tubes (Testim, Vogelxo, generics)

| 7. Do the benefits of continued therapy outweigh the risks? | □ Yes | □ No | |
|---|--|--|--|
| outweigh the risks? | Sign and date on page 3 | STOP | |
| | | Coverage not approved | |
| 8. What is the indication or diagnosis? | ☐ Hypogonadism - Proceed to question 9 | | |
| | ☐ Female-to-male gender dysphoria hormone therapy in a natal female patient (assigned female at birth) - Proceed to question 16 | | |
| | ☐ Other - Proceed to question 24 | | |
| 9. Is the patient a male who is 18 years of age or older? | □ Yes | □ No | |
| C. C | Proceed to question 10 | STOP | |
| | | Coverage not approved | |
| 10. Does the patient have a confirmed diagnosis of hypogonadism as evidenced | □ Yes | □ No | |
| by morning total serum testosterone levels below 300 ng/dL taken on at least two separate occasions? | Proceed to question 12 | Proceed to question 11 | |
| 11. Is testosterone being prescribed by an endocrinologist or urologist who has made | ☐ Yes | □ No | |
| the diagnosis of hypogonadism based on | Proceed to question 12 | STOP | |
| unequivocally and consistently low serum total testosterone or free testosterone levels? | | Coverage not approved | |
| 12. Is the patient experiencing signs and symptoms associated with hypogonadism? | □ Yes | □ No | |
| symptoms associated with hypogoniadism: | Proceed to question 13 | STOP | |
| | | Coverage not approved | |
| 13. Has the provider investigated the etiology of the low testosterone levels? | □ Yes | □ No | |
| of the low testesterone levels. | Proceed to question 14 | STOP | |
| | | Coverage not approved | |
| 14. Has the provider assessed the risks versus benefits of initiating testosterone therapy in | □ Yes | □ No | |
| this patient? | Proceed to question 15 | STOP | |
| | | Coverage not approved | |
| 15. Does the provider acknowledge that testosterone therapy is clinically | □ Yes | □ No | |
| appropriate and needed? | Proceed to question 25 | STOP | |
| | | Coverage not approved | |
| 16. Is the indication for initiation or continuation of female-to-male gender dysphoria hormone therapy in a natal female patient (assigned female at birth)? | ☐ Initiation of female-to-male gender dysphoria hormone therapy in a natal female patient (assigned female at birth)- Proceed to question 17 | | |
| | ☐ Continuation of female-to-male gender dysphoria hormone therapy in a natal female patient (assigned female at birth)-Proceed to question 18 | | |
| 17. Is the patient a female active duty servicemember? | ☐ Yes (Female active duty servicement not approved | emale active duty servicemembers) – STOP - Coverage red | |
| | □ No (Female non-active duty servicemembers) - Proceed to question 18 | | |
| | | | |

USFHP Prior Authorization Request Form for

Androgel, testosterone 1% and 1.62% gel MDP and gel packets, 2% testosterone solution MDP, 1% testosterone gel in unit-dose tubes (Testim, Vogelxo, generics)

| 18. Is the pa | itient 19 years of age or older? | ☐ Yes | □ No | |
|---------------------------------|---|---|------------------------|--|
| | | Proceed to question 19 | STOP | |
| | | | Coverage not approved | |
| | e patient have a diagnosis of | □ Yes | □ No | |
| | dysphoria made by a TRICARE- ed mental health provider | Proceed to question 20 | STOP | |
| accordii DSM? | ng to the most current edition of the | | Coverage not approved | |
| | quested medication being ed by an endocrinologist or a | ☐ Yes | □ No | |
| physicia | physician who specializes in the treatment of transgender patients? | Proceed to question 21 | STOP | |
| of trans | | | Coverage not approved | |
| | ntient a biological female of nring potential? | ☐ Yes | □ No | |
| Ciliubea | ining potential? | Proceed to question 22 | Proceed to question 23 | |
| | | | | |
| 22. Is the pa | tient pregnant or breastfeeding? | ☐ Yes | □ No | |
| | | STOP | Proceed to question 23 | |
| | | Coverage not approved | | |
| 23. Does the | e patient have a psychiatric dity that would confound a | ☐ Yes | □ No | |
| diagnos | is of gender dysphoria or interfere | STOP | Proceed to question 25 | |
| body dy or other | atment (for example: unresolved smorphic disorder; schizophrenia psychotic disorders that have not abilized with treatment)? | Coverage not approved | | |
| | nt the requested indication and e for use. | | | |
| | | | | |
| | | Proceed to question 25 | | |
| 25. What is | the requested medication? | | | |
| | · | ☐ testosterone 1% gel (for example, generic Androgel, generic Testim, etc.) - Sign and date below | | |
| | | ☐ testosterone 1.62% gel (for example, generic Androgel, etc.) - Sign and date below | | |
| | | □ testosterone 2% solution (for example, generic Axiron, etc.) - Sign and date below □ Other (for example Androderm, Fortesta (2% testosterone gel multi-dose pump (MDP)), Natesto, brand Testosterone 1% gel packet, brand Vogelxo) - Proceed to question 26 | | |
| | | | | |
| | Has the patient tried and failed a 3-month trial of one of the following medications: | □ Yes | □ No | |
| testoste Androge (for exa | rone 1% gel (for example, generic el, generic Testim, etc.), 1.62% gel mple, generic Androgel, etc.), or 2% (for example, generic Axiron, etc.)? | Sign and date below | Proceed to question 27 | |

USFHP Prior Authorization Request Form for

Androgel, testosterone 1% and 1.62% gel MDP and gel packets, 2% testosterone solution MDP, 1% testosterone gel in unit-dose tubes (Testim, Vogelxo, generics)

| | 27. Has the patient experienced a clinically significant adverse reaction to one of the following medications: testosterone 1% gel (for example, generic Androgel, generic Testim, etc.), 1.62% gel (for example, generic Androgel, etc.), or 2% solution (for example, generic Axiron, etc.)? | ☐ Yes Sign and date below | ☐ No Proceed to question 28 | | |
|-----------|--|---------------------------|---------------------------------|--|--|
| | 28. Has the patient had a contraindication or relative contraindication to one of the following medications: testosterone 1% gel (for example, generic Androgel, generic Testim, etc.), 1.62% gel (for example, generic Androgel, etc.), or 2% solution (for example, generic Axiron, etc.)? | ☐ Yes Sign and date below | □ No STOP Coverage not approved | | |
| Step 3 | I certify the above is true to the best of my knowledge. Please sign and date: | | | | |
| | Prescriber Signature | Date | | | |
| | | | [26 May 2025] | | |

[26 May 2025]