## US Family Health Plan Prior Authorization Request Form for Adalimumab (**Humira**, **Abbvie Only**)

To be completed and signed by the prescriber. To be used only for prescriptions which are to be filled through the Department of Defense (DoD) US Family Health Plan Pharmacy Program. US Family Health Plan is a TRICARE contractor for DoD.

The completed form may be faxed to 855-273-5735

OR

The patient may attach the completed form to the prescription and **mail** it to: Attn: Pharmacy, 77 Warren St, Brighton, MA 02135

QUESTIONS? Call 1-877-880-7007

When prescribed by a rheumatologist, prior authorization is not required. Prior authorization is required when prescribed in other situations.

Note that the PA applies to the branded Humira formulation by Abbvie. The Cordavis brand PA is on a separate PA form and requires use of Humira first.

Prior authorization does not expire.

Step	Please complete patient and physician information (please print):					
1	Patient Name:Address:		Physician Name:Address:			
					Spapage ID #	
		Sponsor ID # Date of Birth:		Phone #:		
	Secure Fax #:					
Step 2	Please complete the clinical assessment:					
	1. Is the medication being prescribed by a		□ Yes	🗆 No		
	rheumatologist?		Sign and date below	proceed to question 2		
	2. Is the patient 18 years of	age or older?	□ Yes	🗆 No		
			proceed to question 9	proceed to question 3		
	3. What is the indication or	r 🛛 moderate to severe active polyarticular juvenile idiopathic arthritis (pJIA) - proceed				
	diagnosis in this pediatric patient?	to question 4				
	pediatric patient?	<ul> <li>treatment of uveitis (non-infectious intermediate, posterior and panuveitis patients) - proceed to question 4</li> <li>moderately to severely active Crohn's disease – proceed to question 6</li> <li>hidradenitis suppurativa – go to question 7</li> </ul>				
		□ Severe chronic plaque psoriasis in patients who are candidates for systemic or				
		phototherapy, and when other systemic therapies are medically less appropriate (4-17				
		years) – go to question <b>8</b>				
		□ moderately to severely active <b>ulcerative colitis</b> – go to question <b>5</b>				
		<ul> <li>Other indication or diagnosis – STOP: Coverage not approved.</li> <li>Please document diagnosis:</li> </ul>				
	4. Is the patient 2 years of age or older?		□ Yes	🗆 No		
			proceed to question 13	STOP		
				Coverage not approved		

## US Family Health Plan Prior Authorization Request Form for Adalimumab (Humira, Abbvie Only)

			T	
5.	Is the patient 5 years of age or older?		☐ Yes proceed to question <b>8</b>	☐ No STOP Coverage not approved
6.	Is the patient 6 years of age or older?		☐ Yes proceed to question <b>13</b>	☐ No STOP Coverage not approved
7.	Is the patient 12 years of age or older?		☐ Yes proceed to question <b>13</b>	No STOP Coverage not approved
8.	Has the patient had an inadequate response to non- biologic systemic therapy? (For example: methotrexate, aminosalicylates [such as, sulfasalazine, mesalamine], corticosteroids, immunosuppressants [such as, azathioprine], etc.)?		☐ Yes proceed to question <b>13</b>	☐ No STOP Coverage not approved
9.	What is the indication	moderately to severely activ	ve <b>rheumatoid arthritis</b> – go to o	uestion <b>12</b>
	or diagnosis in this	active psoriatic arthritis -		
	adult patient?	🛛 🗆 Ankylosing spondylitis – 🤅	go to question <b>10</b>	
		Active non-radiographic axial spondyloarthritis (nr-ax SpA) with objective signs of inflammation – go to question 12		
		moderate to severe chronic plaque psoriasis in a patient who may benefit from taking injection or pills (systemic therapy) or phototherapy – go to question 12		
		moderately to severely active Crohn's disease – go to question 11		
		moderately to severely active ulcerative colitis – go to question 12		
		moderately to severely activ	e ulcerative colitis – go to ques	Stion 12
		hidradenitis suppurativa -	- go to question <b>13</b>	
		hidradenitis suppurativa -	- · ·	
		<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> </ul>	- go to question <b>13</b> fectious intermediate, posterior a re <b>pyoderma gangrenosum (PC</b>	nd panuveitis patients)– go
		<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids- go</li> </ul>	- go to question <b>13</b> fectious intermediate, posterior a re <b>pyoderma gangrenosum (PC</b>	nd panuveitis patients)– gc 6) that is refractory to high-
		<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids- go</li> <li>Other indication or diagnosis</li> </ul>	- go to question <b>13</b> fectious intermediate, posterior a re <b>pyoderma gangrenosum (PC</b> to question <b>13</b>	nd panuveitis patients)– go 6) that is refractory to high- oved.
10.	Has the patient had an ir	<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids- go</li> <li>Other indication or diagnosis</li> <li>Please document diagnosis:</li> </ul>	- go to question 13 fectious intermediate, posterior a re <b>pyoderma gangrenosum (PC</b> to question 13 s – <b>STOP: Coverage not appr</b>	nd panuveitis patients)– go 6) that is refractory to high- oved.
10.	least two NSAIDS over a	<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids - go</li> <li>Other indication or diagnosis:</li> <li>Please document diagnosis:</li> <li>madequate response to at</li> </ul>	- go to question <b>13</b> fectious intermediate, posterior a re <b>pyoderma gangrenosum (PC</b> to question <b>13</b> s – <b>STOP: Coverage not appr</b>	nd panuveitis patients)– go 6) that is refractory to high- oved.
10.		<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids - go</li> <li>Other indication or diagnosis:</li> <li>Please document diagnosis:</li> <li>madequate response to at</li> </ul>	- go to question 13 fectious intermediate, posterior a re pyoderma gangrenosum (PC to question 13 s – STOP: Coverage not appr	nd panuveitis patients)– go 6) that is refractory to high- oved.
	least two NSAIDS over a months?	<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids- go</li> <li>Other indication or diagnosis:</li> <li>Please document diagnosis:</li> <li>nadequate response to at period of at least two</li> </ul>	- go to question 13 fectious intermediate, posterior a re pyoderma gangrenosum (PC to question 13 s – STOP: Coverage not appr	nd panuveitis patients)– gc b) that is refractory to high- oved. No STOP
	least two NSAIDS over a	<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids- go</li> <li>Other indication or diagnosis:</li> <li>Please document diagnosis:</li> <li>nadequate response to at period of at least two</li> </ul>	- go to question 13 fectious intermediate, posterior a re <b>pyoderma gangrenosum (PC</b> to question 13 s – <b>STOP: Coverage not appr</b> Yes proceed to question 13	nd panuveitis patients)– go b) that is refractory to high- oved. No STOP Coverage not approved No
11.	least two NSAIDS over a months? Does the patient have fis Has the patient had an in biologic systemic therap methotrexate, aminosali sulfasalazine, mesalamin	<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids- go</li> <li>Other indication or diagnosis:</li> <li>Please document diagnosis:</li> <li>Please document diagnosis:</li> <li>adequate response to at period of at least two</li> <li>stulizing CD?</li> </ul>	- go to question 13 fectious intermediate, posterior a re pyoderma gangrenosum (PC to question 13 s – STOP: Coverage not appr Yes proceed to question 13	nd panuveitis patients)– go b) that is refractory to high- oved. No STOP Coverage not approved No proceed to question 12 No STOP
11.	least two NSAIDS over a months? Does the patient have fis Has the patient had an ir biologic systemic therap methotrexate, aminosali sulfasalazine, mesalamin immunosuppressants [s Cases of worsening con and new onset CHF have	<ul> <li>hidradenitis suppurativa -</li> <li>treatment of uveitis (non-into question 12</li> <li>moderately to severely active potency corticosteroids- go</li> <li>Other indication or diagnosis:</li> <li>Please document diagnosis:</li> <li>Please document diagnosis:</li> <li>adequate response to at period of at least two</li> <li>stulizing CD?</li> </ul>	- go to question 13 fectious intermediate, posterior a re pyoderma gangrenosum (PC to question 13 s – STOP: Coverage not appr Ves proceed to question 13 Yes proceed to question 13 Yes	nd panuveitis patients)– gc b) that is refractory to high- oved. No STOP Coverage not approved No proceed to question 12 No

## Adalimumab (Humira, Abbvie Only)

15. Will the patient be receiving other targeted immunomodulatory biologics with Humira, including but not limited to the following: certolizumab (Cimzia), etanercept (Enbrel), golimumab (Simponi), infliximab (Remicade), apremilast (Otezla), ustekinumab (Stelara), abatacept (Orencia), anakinra (Kineret), tocilizumab (Actemra), tofacitinib (Xeljanz/Xeljanz XR), rituximab (Rituxan), secukinumab (Cosentyx), ixekizumab (Taltz), brodalumab (Siliq), sarilumab (Kevzara), guselkumab (Tremfya), baricitinib (Olumiant), tildrakizumab (Ilumya), risankizumab (Skyrizi), or upadacitinib (Rinvoq ER)?	☐ Yes STOP Coverage not approved	☐ No Sign and date below
---	--	-----------------------------

Step	I certify the above is true to the best of my knowledge. Please sign and date:
3	

Prescriber Signature

Date

[17 April 2024]