

USFHP Prior Authorization Request Form for
enlicitide decanoate (Lipfendra)

To be completed and signed by the prescriber. To be used only for prescriptions which are to be filled through the Department of Defense (DoD) US Family Health Plan Pharmacy Program. US Family Health Plan is a TRICARE contractor for DoD.

The completed form may be faxed to 855-273-5735

OR

The patient may attach the completed form to the prescription and **mail** it to:

Attn: Pharmacy, 77 Warren St, Brighton, MA 02135

QUESTIONS? Call 1-877-880-7007

<https://www.usfamilyhealth.org/for-providers/pharmacy-information/>

Prior authorization expires in 1 year. Note that initial TRICARE PA approval is required for renewal. PA will be renewed indefinitely. Clinical documentation may be required.

Step 1 Please complete patient and physician information (please print):

1

Patient Name: _____

Physician Name: _____

Address: _____

Address: _____

Sponsor ID # _____

Phone #: _____

Date of Birth: _____

Secure Fax #: _____

Step 2 Please complete the clinical assessment:

2

1. Has the patient received this medication under the TRICARE benefit in the last 6 months? Please choose "No" if the patient did not previously have a TRICARE approved PA for the requested medication. Note that a trial of Repatha is required first.	☐ Yes (Subject to verification) Proceed to question 27	☐ No Proceed to question 2
2. Provider acknowledges that both Praluent and Repatha are also indicated for the reduction in the risk of major adverse cardiovascular (CV) events in adults at increased risk and have favorable CV outcomes data in adults with established CV disease. At this time, Lipfendra does not have this type of data.	☐ Acknowledged Proceed to question 3	
3. Has the patient tried and failed therapy with evolocumab (Repatha)?	☐ Yes Proceed to question 5	☐ No Proceed to question 4
4. Has the patient experienced a significant adverse reaction to evolocumab (Repatha) that is not expected to occur with the requested medication?	☐ Yes Proceed to question 5	☐ No STOP Coverage not approved
5. Is the patient 18 years of age or older?	☐ Yes Proceed to question 6	☐ No STOP Coverage not approved
6. What is the diagnosis or indication?	☐ Heterozygous familial hypercholesterolemia (HeFH) - Proceed to question 7 ☐ Clinical atherosclerotic cardiovascular disease (ASCVD) - Proceed to question 7 ☐ High risk for ASCVD - Proceed to question 7 ☐ Other – STOP – Coverage not approved	

**USFHP Prior Authorization Request Form for
enlicitide decanoate (Lipfendra)**

7. Will the patient be on concurrent statin therapy at a maximal tolerated dose while on the requested medication?	☐ Yes Proceed to question 12	☐ No Proceed to question 8
8. Has the patient experienced intolerable and persistent (for longer than 2 weeks) muscle symptoms (muscle pain, weakness, cramps) while on statin therapy?	☐ Yes Proceed to question 9	☐ No Proceed to question 11
9. Has the patient undergone at least 2 trials of statin re-challenges with reappearance of muscle symptoms? -- NOTE: that is, the patient has had 2 trials of statins with muscle symptoms.	☐ Yes Proceed to question 12	☐ No Proceed to question 10
10. Has the patient had a creatine kinase (CK) level greater than 10 times the upper limit of normal OR rhabdomyolysis with CK greater than 10,000 international units per liter (IU/L) that is unrelated to statin use?	☐ Yes Proceed to question 12	☐ No Proceed to question 11
11. Does the patient have a contraindication to the use of a statin? -- NOTE: Please select the option that best applies to this patient's condition.	☐ Active Liver Disease (including unexplained persistent elevations in hepatic transaminase levels) - Proceed to question 12 ☐ Hypersensitivity - Proceed to question 12 ☐ Pregnancy - Proceed to question 12 ☐ Nursing mothers - Proceed to question 12 ☐ None of the above - STOP – Coverage not approved	
12. Is the patient pregnant or breastfeeding?	☐ Yes STOP Coverage not approved	☐ No Proceed to question 13
13. What is the indication or diagnosis?	☐ Heterozygous familial hypercholesterolemia (HeFH) – Sign and date below ☐ Clinical atherosclerotic cardiovascular disease (ASCVD) - Proceed to question 14 ☐ High risk for ASCVD - Proceed to question 20 ☐ Other – STOP – Coverage not approved	
14. Is the patient at very high risk for future ASCVD events? NOTE: patients at very high risk for future ASCVD events include those with a history of multiple major ASCVD events or 1 major ASCVD event and multiple high-risk conditions. Refer to the 2022 ACC Expert Consensus Decision Pathway on the role of non-statin therapies for LDL-cholesterol lowering in the management of ASCVD for more information.	☐ Yes Proceed to question 15	☐ No Proceed to question 16
15. Does the patient have an LDL level greater than 55 mg/dL despite statin at maximal tolerated doses?	☐ Yes Proceed to question 17	☐ No STOP Coverage not approved
16. Does the patient have an LDL level greater than 70 mg/dL despite statin at maximal tolerated doses?	☐ Yes Proceed to question 17	☐ No STOP Coverage not approved
17. Has the patient tried EITHER atorvastatin (Lipitor) at a dose of 40 mg to 80 mg OR rosuvastatin (Crestor) at a dose of 20 mg to 40 mg for at least 4 to 6 weeks each?	☐ Yes Sign and date below	☐ No Proceed to question 18

**USFHP Prior Authorization Request Form for
enlicitide decanoate (Lipfendra)**

18. Has the patient tried any statin at a maximally tolerated dose in combination with ezetimibe (Zetia) for at least 4 to 6 weeks?	☐ Yes Sign and date below	☐ No Proceed to question 19
19. Has the patient tried ezetimibe (Zetia) as monotherapy (alone) for at least 4 to 6 weeks?	☐ Yes Sign and date below	☐ No STOP Coverage not approved
20. Does the patient have LDL level greater than 190 mg/dL?	☐ Yes Proceed to question 24	☐ No Proceed to question 21
21. Does the patient have diabetes and LDL level less than 190 mg/dL?	☐ Yes Proceed to question 24	☐ No Proceed to question 22
22. Does the patient have LDL 70 to 189 mg/dL and an estimated 10-year risk for ASCVD greater than 7.5%?	☐ Yes Proceed to question 24	☐ No Proceed to question 23
23. Does the patient have LDL level less than 190 mg/dL and evidence of significant subclinical atherosclerosis defined as: Significant atherosclerotic plaque observed in an asymptomatic patient on any of the following diagnostic studies: coronary artery calcification noted on computed tomography (CT) studies, including calcium scoring, cardiac CT coronary angiography, chest CT for ruling out pulmonary embolism, chest CT for lung cancer screening, or diagnostic chest CT; carotid plaque noted on carotid ultrasound or angiography; or abnormal ankle-brachial index or plaque noted on peripheral arterial angiography?	☐ Yes Proceed to question 24	☐ No STOP Coverage not approved
24. Has the patient tried EITHER atorvastatin (Lipitor) at a dose of 40 mg to 80 mg OR rosuvastatin (Crestor) at a dose of 20 mg to 40 mg for at least 4 to 6 weeks each?	☐ Yes Sign and date below	☐ No Proceed to question 25
25. Has the patient tried any statin at a maximally tolerated dose in combination with ezetimibe (Zetia) for at least 4 to 6 weeks?	☐ Yes Sign and date below	☐ No Proceed to question 26
26. Is the patient statin intolerant, and has tried ezetimibe (Zetia) as monotherapy (alone) for at least 4 to 6 weeks?	☐ Yes Sign and date below	☐ No STOP Coverage not approved
27. Has the patient responded to therapy, as evidenced by at least a 50% reduction in LDL levels from baseline?	☐ Yes Sign and date below	☐ No STOP Coverage not approved

Step 3 I certify the above is true to the best of my knowledge. Please sign and date:

Prescriber Signature

Date

[04 August 2026]